

## Portland Harbor Food Web Model Internal Parameters not set at Arnot and Gobas

These are the 21 internal FWM parameters LWG calibrated from Monte Carlo runs. None of these parameters have Portland Harbor site specific information available. A concern of several reviewers was why were model defaults being changed in the table below. The table below lists the FWM parameter, its default Arnot and Gobas model value. Current FWM values are from a spreadsheet sent to Burt Shephard by Elizabeth Al. Most of the current FWM values are on the 'inputs' tab of the 11/4/2014 spreadsheet.

No.	Model parameter	AQUAWEB v1.2 default
<b><i>General biological parameters</i></b>		
	Resistance to chemical uptake through aqueous	
1	phase for phytoplankton (UA)	0.00006
	Resistance to chemical uptake through organic	
2	phase for phytoplankton (UB)	5.5
3	NLOM*-octanol proportionality constant (Beta)	0.035
<b><i>Species specific biological parameters</i></b>		
<u><i>Zooplankton</i></u>		
4	Dietary absorption efficiency of lipid	0.75
5	Dietary absorption efficiency of NLOM	0.75
<u><i>Benthic invertebrate filter feeder (clams)</i></u>		
6	Dietary absorption efficiency of lipid	0.75
7	Dietary absorption efficiency of NLOM	0.75
<u><i>Benthic invertebrate consumers</i></u>		
8	Dietary absorption efficiency of lipid	0.75
9	Dietary absorption efficiency of NLOM	0.75
<u><i>Epibenthic invertebrate consumers (crayfish)</i></u>		
10	Dietary absorption efficiency of lipid	0.75
11	Dietary absorption efficiency of NLOM	0.75
<u><i>Sculpin</i></u>		
12	Dietary absorption efficiency of lipid	0.92
13	Dietary absorption efficiency of NLOM	0.60
<u><i>Largescale sucker</i></u>		
14	Dietary absorption efficiency of lipid	0.92
15	Dietary absorption efficiency of NLOM	0.60
<u><i>Carp</i></u>		
16	Dietary absorption efficiency of lipid	0.92
17	Dietary absorption efficiency of NLOM	0.60
<u><i>Smallmouth bass</i></u>		
18	Dietary absorption efficiency of lipid	0.92
19	Dietary absorption efficiency of NLOM	0.60
<u><i>Northern pikeminnow</i></u>		
20	Dietary absorption efficiency of lipid	0.92
21	Dietary absorption efficiency of NLOM	0.60

\* NLOM = non-lipid organic matter (e.g. proteins, nucleic acids)

## is (2004) default values

instead of using Gobas defaults

for them

absence of site- or species-specific information

from AQUAWEB v1.2, and the current FWM values

len on 11/4/2014

meet from Elizabeth Allen. Must unhide some rows to find all of the current FWM values

Current Portland Harbor FWM value	Units	Comment
0.00006	Unitless	
5.5	Unitless	
0.035	L/kg	
0.72	Proportion	Parameter called alpha or default lipid absorp
0.72	Proportion	Parameter called beta or default NLOC/NLOM
0.75	Proportion	Parameter called alpha or default lipid absorp
0.75	Proportion	Parameter called beta or default NLOC/NLOM
0.75	Proportion	Parameter called alpha or default lipid absorp
0.75	Proportion	Parameter called beta or default NLOC/NLOM
0.75	Proportion	Parameter called alpha or default lipid absorp
0.75	Proportion	Parameter called beta or default NLOC/NLOM
0.92	Proportion	Parameter called alpha or default lipid absorp
0.60	Proportion	Parameter called beta or default NLOC/NLOM
0.92	Proportion	Parameter called alpha or default lipid absorp
0.60	Proportion	Parameter called beta or default NLOC/NLOM
0.92	Proportion	Parameter called alpha or default lipid absorp
0.60	Proportion	Parameter called beta or default NLOC/NLOM
0.92	Proportion	Smallmouth bass is only fish species default A
0.60	Proportion	Smallmouth bass is only fish species default A
0.92	Proportion	Parameter called alpha or default lipid absorp
0.60	Proportion	Parameter called beta or default NLOC/NLOM



tion efficiency in AQUAWEB v1.2. Given as a percentage in Gobas model (i.e. 75%)  
l absorption efficiency in AQUAWEB v1.2. Given as a percentage in Gobas model (i.e. 75%)

tion efficiency in AQUAWEB v1.2. Given as a percentage in Gobas model (i.e. 75%)  
l absorption efficiency in AQUAWEB v1.2. Given as a percentage in Gobas model (i.e. 75%)

tion efficiency in AQUAWEB v1.2. Given as a percentage in Gobas model (i.e. 75%)  
l absorption efficiency in AQUAWEB v1.2. Given as a percentage in Gobas model (i.e. 75%)

tion efficiency in AQUAWEB v1.2. Given as a percentage in Gobas model (i.e. 75%)  
l absorption efficiency in AQUAWEB v1.2. Given as a percentage in Gobas model (i.e. 75%)

tion efficiency in AQUAWEB v1.2. Given as a percentage in Gobas model (i.e. 92%)  
l absorption efficiency in AQUAWEB v1.2. Given as a percentage in Gobas model (i.e. 60%)

tion efficiency in AQUAWEB v1.2. Given as a percentage in Gobas model (i.e. 92%)  
l absorption efficiency in AQUAWEB v1.2. Given as a percentage in Gobas model (i.e. 60%)

tion efficiency in AQUAWEB v1.2. Given as a percentage in Gobas model (i.e. 92%)  
l absorption efficiency in AQUAWEB v1.2. Given as a percentage in Gobas model (i.e. 60%)

QUAWEB v1.2 has in common with Portland Harbor FWM. Called Fish 12 in AQUAWEB. Parameter call  
QUAWEB v1.2 has in common with Portland Harbor FWM. Called Fish 12 in AQUAWEB. Parameter call

tion efficiency in AQUAWEB v1.2. Given as a percentage in Gobas model (i.e. 92%)  
l absorption efficiency in AQUAWEB v1.2. Given as a percentage in Gobas model (i.e. 60%)



ed alpha or default lipid absorption efficiency in AQUAWEB v1.2. Given as a percentage in Gobas model  
ed beta or default NLOC/NLOM absorption efficiency in AQUAWEB v1.2. Given as a percentage in Goba:





(i.e. 90%)  
s model (i.e. 60%)